

The information above this line is not included in the label

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



RLP
Approved



(Dog photograph wrapping to carton sides)

advantix[®]

For Puppies and Small Dogs up to 4 kg

100 g/L IMIDACLOPRID
500 g/L PERMETHRIN (40:60)
Solvent: 484 g/L N-METHYLPYRROLIDONE

Fleas & ticks

[PLACE DOG AND CAT SYMBOLS]
Width 20 mm Height 18 mm
Text 1.9 mm and 1.5 mm – (Keyline version)

Once a month topical flea and tick treatment

- Apply fortnightly for paralysis ticks
- Kills and repels sandflies, mosquitoes and ticks
- Stops fleas feeding in 3-5 minutes

[Water resistant formulation (*text in pictogram*)]

[1 x 0.4 mL] [2 x 0.4 mL] [3 x 0.4 mL] [6 x 0.4 mL]
(*tube pictogram*)

(*tick pictogram*) (*flea pictogram*) (*larvae pictogram*) (*mosquito pictogram*) (*louse pictogram*) (*stable fly pictogram*)

Carton back panel	Issue Date: 9Jan2013 Version: 3	Page: 2 of 10
The information above this line is not included in the label		

advantix[®]

For Puppies and Small Dogs up to 4 kg

For:

- the control of ticks including the paralysis tick *Ixodes holocyclus*, brown dog tick *Rhipicephalus sanguineus*, and bush tick *Haemaphysalis longicornis*. ADVANTIX both repels and kills ticks;
- the treatment and prevention of fleas *Ctenocephalides* spp. Reduces the incidence of Flea Allergy Dermatitis. Larval flea stages are killed in the surroundings of ADVANTIX treated pets;
- the control of lice *Trichodectes canis*, *Linognathus setosus* for up to 6 weeks;
- the repelling and killing of mosquitoes (*Anopheles*, *Culex*, *Aedes* sp) and sandflies (*Phlebotomus*, *Culicoides* sp);
- the repelling of stable (biting) flies (*Stomoxys calcitrans*) in small dogs and puppies.

One application lasts for a month; or 2 weeks for paralysis tick.

Read the enclosed leaflet before using.

DIRECTIONS FOR USE

[PLACE DOG AND CAT SYMBOLS]

Width 20 mm Height 18 mm

Text 1.9 mm and 1.5 mm

Keep cats separated from recently treated dogs.

There are reports of toxic effect in cats which groom or contact dogs treated 48 hours earlier

Toxic effects include behaviour changes, drooling, tremors and death

Seek veterinary advice immediately if you suspect toxicity in a cat.

For external use only.

One tube treats small dogs and puppies up to 4 kg bodyweight.

During tick season daily searching for and removal of any ticks found is recommended.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

SAFETY DIRECTIONS

Will damage the eyes. Avoid contact with the eyes. If product in eyes, wash it out immediately with water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

DISPOSAL

After use replace cap on tube. Dispose of used applicators by wrapping in paper and placing in garbage

Carton back panel (cont'd)	Issue Date 9Jan2013 Version: 3	Page: 3 of 10
<i>The information above this line is not included in the label</i>		

STORAGE

Store below 30°C (Room Temperature)

For batch and expiry information see bottom of carton

APVMA Approval No: 58259/57498

Bayer

Bayer Australia Ltd
875 Pacific Highway
PYMBLE NSW 2073
ACN 000 138 714

Limitation of Liability

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, BAL's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

ADVANTIX® is a registered trademark of Bayer AG, Leverkusen, Germany.

Manufactured in Germany

08/12

Carton side panel 1	Issue Date 9Jan2013 Version: 3	Page: 4 of 10
<i>The information above this line is not included in the label</i>		

advantix® for puppies and small dogs up to 4 kg

[1 x 0.4 mL] [2 x 0.4 mL] [3 x 0.4 mL] [6 x 0.4 mL]

Fleas & ticks

- Kills and repels sandflies, mosquitoes and ticks
- Stops fleas feeding in 3-5 minutes

Carton side panel 2	Issue Date 9Jan2013 Version: 3	
<i>The information above this line is not included in the label</i>		

advantix® for puppies and small dogs up to 4 kg

[1 x 0.4 mL] [2 x 0.4 mL] [3 x 0.4 mL] [6 x 0.4 mL]

Fleas & ticks

CUSTOMER INFORMATION LINE

1800 678 368 TOLL FREE from anywhere in Australia For a free
treatment reminder visit bayeranimal.com.au [1, 2, 3, 6 x 0.4 mL]

9.00am to 4.00pm (Eastern Time) Monday to Friday

friendsforever.com.au

Carton top flap	Issue Date 9Jan2013 Version: 3	
<i>The information above this line is not included in the label</i>		

advantix® for puppies and small dogs up to 4 kg

[1 x 0.4 mL] [2 x 0.4 mL] [3 x 0.4 mL] [6 x 0.4 mL]

Fleas & ticks

Carton bottom flap	Issue Date 9Jan2013 Version: 3	
<i>The information above this line is not included in the label</i>		

Batch

Expiry

Foil Pouch label	Issue Date 9Jan2013 Version: 3	Page: 5 of 10
<i>The information above this line is not included in the label</i>		

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



advantix[®]

For Puppies and Small Dogs up to 4 kg

100 g/L IMIDACLOPRID

500 g/L PERMETHRIN (40:60)

Solvent: 484 g/L N-METHYLPYRROLIDONE

Read accompanying instructions before use

Batch:

APVMA 58259/57498

Exp:

[PLACE DOG AND CAT SYMBOLS]

Width 20 mm Height 18 mm

Text 1.9 mm and 1.5 mm

THIS PRODUCT IS LABELLED FOR SALE IN AUSTRALIA ONLY.
IT IS ILLEGAL TO SELL THIS PRODUCT – AS LABELLED – IN THE USA.
[1, 2, 3, 6 x 0.4 mL]

Common leaflet	Issue Date 9Jan2013 Version: 3	Page: 6 of 10
The information above this line is not included in the label		

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



advantix®

For Puppies and Small Dogs up to 4 kg

100 g/L IMIDACLOPRID
500 g/L PERMETHRIN (cis:trans, 40:60)
Solvent: 484 g/L N-METHYLPYRROLIDONE

For:

- the control of ticks including the paralysis tick *Ixodes holocyclus*, brown dog tick *Rhipicephalus sanguineus*, and bush tick *Haemaphysalis longicornis*. ADVANTIX both repels and kills ticks;
- the treatment and prevention of fleas *Ctenocephalides* spp. Reduces the incidence of Flea Allergy Dermatitis. Larval flea stages are killed in the surroundings of ADVANTIX treated dogs;
- for the control of lice *Trichodectes canis*, *Linognathus setosus* for up to 6 weeks;
- the repelling and killing of mosquitoes (*Anopheles*, *Culex*, *Aedes* sp) and biting midges/sandflies (*Phlebotomus*, *Culicoides* sp);
- the repelling of stable (biting) flies (*Stomoxys calcitrans*) in dogs.

One application lasts for a month, (2 weeks for paralysis tick).

DIRECTIONS FOR USE

For external application only.

[PLACE DOG AND CAT SYMBOLS]

Width 20 mm Height 18 mm

Text 1.9 mm and 1.5 mm

Keep cats separated from recently treated dogs

There are reports of toxic effect in cats which groom or contact dogs treated 48 hours earlier

Toxic effect include behaviour changes, drooling, tremors and death

Seek veterinary advice immediately if you suspect toxicity in a cat.

Apply every 2 weeks for control of paralysis ticks (*Ixodes holocyclus*). Apply monthly for control of all other ticks and/or fleas.

DOSAGE

For animals of this type ↓		Use 1 TUBE of the colour ↓
Small dogs & puppies	Up to 4 kg	green
Dogs	4 - 10 kg	turquoise
Dogs	10 – 25 kg	pink
Dogs	25 – 50 kg	blue

For very large dogs over 50 kg, apply 2 tubes of ADVANTIX for Dogs Over 25 kg (blue) by dividing the dose between the shoulder blades, on the midline of the back between the hips, and at one point in between.

1. Remove required tube(s) from the package
2. Hold tube upright and pull off the cap.
3. Reverse the cap and push cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Site of Application

graphic
graphic
graphic

Dogs under 25 kg	Dogs 25 - 50 kg
1 TUBE Apply between the shoulder blades	1 TUBE Apply by dividing the dose between the shoulder blades, on the midline of the back between the hips, and at one point in between
<i>Treatment graphic</i>	<i>Treatment graphic</i>

6. The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the tube on the skin and squeeze the tube firmly **twice to apply the solution directly on the skin.** Avoid pet's eyes or mouth.
7. Record treatment date and repeat monthly or fortnightly as directed above. ADVANTIX remains efficacious following soap free shampoo treatment, swimming or exposure to rain. If washing animal during use of this product use a soap free shampoo 14 days after treatment..

GENERAL INSTRUCTIONS
TICK CONTROL

- ADVANTIX must be applied at 2 week intervals for control of paralysis ticks (*Ixodes holocyclus*)
- Application at one month intervals will control brown dog ticks and bush ticks (*Rhipicephalus* sp., *Haemaphysalis* sp.)

Common leaflet (cont'd)	Issue Date 9Jan2013 Version: 3	Page: 8 of 10
<i>The information above this line is not included in the label</i>		

- Ticks are repelled and/or killed
- Ideally use Advantix 1 month prior to first expected exposure to ticks
- Attached ticks will be killed but remain attached and visible
- The Paralysis Tick (*Ixodes holocyclus*) is a potentially deadly parasite and difficult to control
- While Advantix controls paralysis ticks for 2 weeks, the risk of tick paralysis cannot be entirely eliminated. Therefore daily searching for and removal of any ticks found is recommended.

FLEA CONTROL AND PREVENTION

ADVANTIX has been shown to kill adult and larval fleas within 20 minutes of contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite the animal to receive a lethal dose.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with ADVANTIX rapidly kills fleas and reduces the incidence of this condition.

ADVANTIX may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the dog. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product. It is recommended that all dogs in the same environment should be protected by ADVANTIX for Dogs and all cats with ADVANTAGE or ADVOCATE for Cats.

Do not use ADVANTIX on cats.

MOSQUITOES and MIDGES/SANDFLIES

One treatment of ADVANTIX provides repellent and killing activity against mosquitoes and biting midges/sandflies for four weeks thus reducing irritation and the risk of disease transmission.

STABLE (BITING) FLIES

One treatment of ADVANTIX provides repellent activity against stable (biting) flies for four weeks which helps reduce the irritation and skin lesions caused by these pests.

PUPPIES

Puppies may be treated with ADVANTIX from 7 weeks of age. If puppies younger than 7 weeks require flea control they may be safely treated with ADVANTAGE® for Dogs.

PREGNANCY AND LACTATION

ADVANTIX is safe for use on pregnant and lactating bitches.

PRECAUTIONS

DO NOT apply to cats or animals other than dogs. Due to their unique physiology and inability to metabolise certain compounds, the product must not be used on cats. If applied to a cat, or ingested by a cat, this product may have serious harmful effects. If this occurs consult your veterinary surgeon immediately.

Consult your veterinary surgeon before use on sick and debilitated dogs.

Take care to ensure the product does not contact the eyes or mouth of recipient dogs.

The information above this line is not included in the label

On rare occasions reactions on dogs may include skin sensitivity resulting in symptoms such as scratching, agitation or lethargy. These signs are normally transient. If you are concerned please contact Bayer for further information.

ADVANTIX has no strong chemical smell. The liquid vehicle in ADVANTIX may discolour certain surfaces and fabrics e.g., polished floorboards, leather. Exclude the animal from contact with such surfaces until the product dries on the animal's coat.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

SAFETY DIRECTIONS

Will damage the eyes. Avoid contact with the eyes. If product in eyes, wash it out immediately with water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

DISPOSAL

After use replace cap on tube. Dispose of used tubes and packaging by wrapping in paper and placing in garbage.

PRESENTATION

ADVANTIX for	Tube Colour	Contents	APVMA Approval No:
Puppies & Small Dogs up to 4 kg	green	0.4 mL	58259/57498
Dogs 4 – 10 kg	turquoise	1.0 mL	58261/57499
Dogs 10 – 25 kg	pink	2.5 mL	58260/57500
Dogs over 25 kg	blue	4.0 mL	58262/57501

STORAGE

Store below 30°C (Room Temperature)

All pipettes should be used within 18 months of opening the foil pouch

Limitation of Liability

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

Bayer

Bayer Australia Ltd
875 Pacific Highway
PYMBLE NSW 2073
ACN 000 138 714

ADVANTIX[®], ADVOCATE[®] and ADVANTAGE[®] are registered trademarks of Bayer AG, Leverkusen, Germany.
Manufactured in Germany
(art codes)
08/12

Tube label	Issue Date 9Jan2013 Version: 3	Page: 10 of 10
<i>The information above this line is not included in the label</i>		

CAUTION
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



advantix[®]

For Puppies and Small Dogs up to 4 kg

100 g/L IMIDACLOPRID
500 g/L PERMETHRIN
Solvent: 484 g/L N-METHYLPYRROLIDONE

Toxic to cats [font size 1.5 mm high]

0.4 mL



0.4 mL APVMA 58259/57498

Batch:

Exp: