

**<FRONT PANEL>**

**POISON  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING  
OR USING  
FOR ANIMAL TREATMENT ONLY**

**SYKES  
BIG L  
POUR-ON**

**Active constituent: 200 g/L Levamisole (as Levamisole base)**



**For the treatment of levamisole sensitive gastro-intestinal  
parasites and lungworm in cattle.**

**1L  
{2.5L, 5L, 10L, 20L}**

<BACK PANEL>

**Indications:**

**BIG L POUR-ON contains levamisole and is effective against sensitive strains of the following parasites in cattle:**

*(Haemonchus placei)* Barbers pole worm  
*(Ostertagia ostertagi)* Small brown stomach worm (mature adults only) in cattle under 12 months of age\*  
*(Trichostrongylus axei)* Stomach hair worm  
*(Trichostrongylus spp)* Black scour worm  
*(Bunostomum phlebotomum)* Hookworm  
*(Cooperia spp)* Small intestinal worm  
*(Nematodirus spp)* Thin necked intestinal worm  
*(Oesophagostomum radiatum)* Nodule worm  
*(Dictyocaulus viviparus)* Lungworm

\*Where *Ostertagia* is a problem in cattle over 12 months old, products containing benzimidazoles or avermectins/milbemycins are recommended.

**Resistance may develop to any chemical.**

**DIRECTIONS FOR USE:**

**Restraints:**

**DO NOT USE** in lactating cows where milk or milk products may be used for human consumption.

**DO NOT** treat wet animals.

**DO NOT** use oral drenching guns

**Contraindications:**

**This product should not be used on dogs or horses.**

Read directions carefully

**FOR USE ON CATTLE ONLY**

**Dosage and Administration:**

**SHAKE WELL BEFORE USE.**

Calibrate applicator gun before use to ensure accurate dosage. After use, clean gun by flushing out with clean cold water. Do not use hot water.

**PRECAUTIONS**

Normal care should be taken with weak or pregnant animals. Overdose may cause animals to show temporary excitement, salivation or muscular tremors. BIG L POUR-ON may be used

concurrently with pour-on products for lice control provided the products are applied to different parts of the backline.

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.

Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.

Recommended dose should not be exceeded.

### **APPLICATION RATE**

Apply 2.5mL per 50kg live bodyweight evenly along the backline using the appropriate applicator gun. For accurate application, nozzle should be in contact with the skin.

Cattle weight (kg)	Dose volume (mL)	No. of animals treated per 1L/2.5L/ 5L/ 10L/ 20L pack
Up to 50	2.5	400/1000/ 2000/ 4000/ 8000
51-100	5.0	200/500/ 1000/ 2000/ 4000
101-150	7.5	133/333 /666 /1332 /2664
151-200	10.0	100/250/ 500/ 1000/ 2000
201-250	12.5	80/200/ 400/ 800/ 1600
251-300	15.0	66/166/ 332/ 664/ 1328
301-350	17.5	57/142/ 284/ 568/ 1136
351-400	20.0	50/125/ 250/ 500/ 1000
401-450	22.5	44/111/ 222/ 444/ 888
451-500	25.0	40/100/ 200/ 400/ 800
501-550	27.5	36/90/ 180/ 360/ 720
551-600	30.0	33/83/ 166/ 332/ 664
601-650	32.5	30/77/ 154/ 308/ 616

Cattle heavier than 650kg to be dosed at 2.5mL per 50kg bodyweight.

Always dose on a bodyweight basis.

A representative sample of animals should be weighed before treatment either with scales or with a weighband.

The dose rate to be based on the heaviest animal by bodyweight in each mob (cows, steers, calves, bulls).

Do not overdose.

Where there is large variation in size within the group, draft into 2 or more lines based on bodyweight, to avoid excessive overdosing.

Version 1.7 01.10.10 – Big L Pour On - 1L,2.5L,5L,10L,20L- immediate text label  
Size: 116 x 80mm

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 3 days before slaughter for human consumption.**

**MILK: DO NOT USE in lactating cows where milk or milk products may be used for human consumption.**

**TRADE ADVICE: Export Slaughter Interval (ESI):**

**DO NOT USE less than 7 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from either the manufacturer on +61 03 9706 4913 or the APVMA website [www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)**

**SAFETY DIRECTIONS:**

**Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves.**

**FIRST AID:**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.**

**SYKES VET INTERNATIONAL PTY LTD**

**27 Healey Road  
Dandenong VIC 3175**

**DISPOSAL:**

Triple rinse containers into medicated water. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers should not be burned.

**Storage:** Store below 30°C (Room Temperature), in original container. Keep tightly closed in a dry place away from light.

**BATCH No. \_\_\_\_\_**

**EXPIRY DATE \_\_\_\_\_**

**APVMA APPROVAL No. 56795/49123**

**<FRONT PANEL>**

**POISON  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING  
OR USING  
FOR ANIMAL TREATMENT ONLY**

**SYKES  
BIG L  
POUR-ON**

Active constituent: 200 g/L Levamisole (as Levamisole base)



For the treatment of levamisole sensitive gastro-intestinal  
parasites and lungworm in cattle.

125mL  
{250mL, 500mL}

<BACK PANEL>

**Indications:**

**BIG L POUR-ON contains levamisole and is effective against sensitive strains of the following parasites in cattle:**

- (Haemonchus placei)* Barbers pole worm
- (Ostertagia ostertagi)* Small brown stomach worm (mature adults only) in cattle under 12 months of age\*
- (Trichostrongylus axei)* Stomach hair worm
- (Trichostrongylus spp)* Black scour worm
- (Bunostomum phlebotomum)* Hookworm
- (Cooperia spp)* Small intestinal worm
- (Nematodirus spp)* Thin necked intestinal worm
- (Oesophagostomum radiatum)* Nodule worm
- (Dictyocaulus viviparus)* Lungworm

\*Where *Ostertagia* is a problem in cattle over 12 months old, products containing benzimidazoles or avermectins/milbemycins are recommended.

**Resistance may develop to any drench.**

**DIRECTIONS FOR USE:**

**Restraints:**

**DO NOT USE** in lactating cows where milk or milk products may be used for human consumption.

**DO NOT** treat wet animals.

**DO NOT** use oral drenching guns

**Contraindications:**

**This product should not be used on dogs or horses.**

Read directions carefully

**FOR USE ON CATTLE ONLY**

**Dosage and Administration:**

**SHAKE WELL BEFORE USE.**

Calibrate applicator gun before use to ensure accurate dosage. After use, clean gun by flushing out with clean cold water. Do not use hot water.

**PRECAUTIONS**

Normal care should be taken with weak or pregnant animals. Overdose may cause animals to show temporary excitement, salivation or muscular tremors. BIG L POUR-ON may be used concurrently with

pour-on products for lice control provided the products are applied to different parts of the backline.

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.

Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.

Recommended dose should not be exceeded.

### **APPLICATION RATE**

Apply 2.5mL per 50kg live bodyweight evenly along the backline using the appropriate applicator gun. For accurate application, nozzle should be in contact with the skin.

Cattle weight (kg)	Dose volume (mL)	No. of animals treated per 125 mL/250mL/500mL pack
Up to 50	2.5	50/100/200
51-100	5.0	25/50/100
101-150	7.5	16.5/33/66
151-200	10.0	12.5/25/50
201-250	12.5	10/20/40
251-300	15.0	8.5/17/34
301-350	17.5	7/14/28
351-400	20.0	6/12/24
401-450	22.5	5.5/11/22
451-500	25.0	5/10/20
501-550	27.5	4.5/9/18
551-600	30.0	4/8/16
601-650	32.5	3.5/7/14

Cattle heavier than 650kg to be dosed at 2.5mL per 50kg bodyweight. Always dose on a bodyweight basis.

A representative sample of animals should be weighed before treatment either with scales or with a weighband.

The dose rate to be based on the heaviest animal by bodyweight in each mob (cows, steers, calves, bulls).

Do not overdose.

Where there is large variation in size within the group, draft into 2 or more lines based on bodyweight, to avoid excessive overdosing.

Version 1.7 01/10/10 – Big L Pour On - 125mL,250mL,500mL – immediate text label  
Size: 116 x 80mm

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 3 days before slaughter for human consumption.**

**MILK: DO NOT USE in lactating cows where milk or milk products may be used for human consumption.**

**TRADE ADVICE: Export Slaughter Interval (ESI):**

**DO NOT USE less than 7 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from either the manufacturer on +61 03 9706 4913 or the APVMA website [www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)**

**SAFETY DIRECTIONS:**

**Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves.**

**FIRST AID:**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.**

**SYKES VET INTERNATIONAL PTY LTD  
27 Healey Road  
Dandenong VIC 3175**

**DISPOSAL:**

**Dispose of empty container by wrapping with paper and putting in garbage.**

**Storage: Store below 30°C (Room Temperature), in original container. Keep tightly closed in a dry place away from light.**

**BATCH No. \_\_\_\_\_**

**EXPIRY DATE \_\_\_\_\_**

**APVMA APPROVAL No. 56795/49123**